

TUM-HEP-332/98

MPI-PhT/98-86

hep-ph/9811470

October 1998

Fermion mixing renormalization and gauge invariance

P. Gambino^a, P.A. Grassi^b, and F. Madricardo^b

^a *Technische Universität München,
Physik Dept., D-85748 Garching, Germany*

^b *Max Planck Institut für Physik (Werner-Heisenberg-Institut),
Föhringer Ring 6, D-80805 München, Germany*

Abstract

We study the renormalization of the fermion mixing matrix in the Standard Model and derive the constraints that must be satisfied to respect gauge invariance to all orders. We demonstrate that the prescription based on the *on-shell* renormalization conditions is not consistent with the Ward-Takahashi Identities and leads to gauge dependent physical amplitudes. A simple scheme is proposed that satisfies all theoretical requirements and is very convenient for practical calculations.

Despite a few interesting papers have been devoted to the fermion mixing renormalization [1, 2, 3, 4], the subject has so far escaped much attention. This is mostly due to the fact that, as a result of GIM cancelations, the radiative corrections related to the renormalization of the CKM matrix can be made very small, $O(G_F m_q^2)$, where m_q is the mass of a light quark. They are therefore of little practical importance in the context of the Standard Model (SM). Still, the subject has some conceptual interest in its own, not to mention the relevance of mixing in many extension of the SM. In this letter, we reconsider it from a different point of view, which allows us to point out some inconsistency in previous analyses and to propose an alternative solution.

For definiteness, we concentrate on the case of the CKM matrix renormalization in the SM. A convenient framework to study this issue is provided by the Ward Takahashi Identities (WTI) of the theory with background fields [5, 6, 7]. Indeed, as the diagonalization of the fermion mass matrix is achieved by field redefinitions that do not commute with the gauge transformations, the CKM elements appear explicitly in the WTI, unlike masses and gauge couplings. This will give us a strong constraint. At the functional level, the WTI which represent the σ^+ generator of $SU(2)_L$ are implemented by the local functional operator $\mathcal{W}_+$ acting on the effective action Γ (see [7]). For our purposes, the relevant part of $\mathcal{W}_+$ is the one which contains the quark fields:

$$\mathcal{W}_+^{quark} = \sum_{u,d} \left[\bar{\psi}_u^L V_{ud}^0 \frac{\vec{\delta}}{\delta \psi_d^L} - \frac{\overleftarrow{\delta}}{\delta \psi_u^L} V_{ud}^0 \psi_d^L \right]. \quad (1)$$

Here V^0 is the tree level CKM matrix. Upon renormalization, the fermionic fields are rescaled by non-diagonal complex wave function renormalization (WFR) matrices Z_u and Z_d ; consequently, V^0 is replaced by a renormalized CKM matrix $V = (Z_u^L)^{-\frac{1}{2}} V^0 (Z_d^L)^{\frac{1}{2}}$, where Z^L is the left-handed component of the WFR. Expanding V at first order, we obtain for the CKM counterterm

$$\delta V_{ud} = \frac{1}{2} \sum_{u'} \delta Z_{uu'}^L V_{u'd} - \frac{1}{2} \sum_{d'} V_{ud'} \delta Z_{d'd}^L. \quad (2)$$

An additional constraint on the WFR comes from the requirement of unitarity for V ,

$$\sum_{d'} V_{ud'} (\delta Z_{d'd}^L + \delta Z_{dd'}^{L*}) = \sum_{u'} (\delta Z_{uu'}^L + \delta Z_{u'u}^{L*}) V_{u'd}. \quad (3)$$

Notice that this constraint follows also from the request that the commutation relations among the operators $\mathcal{W}_+$, $\mathcal{W}_-$, and $\mathcal{W}_3$ be preserved. Combining Eqs. (2) and (3) we find

$$\delta V_{ud} = -\frac{1}{4} \left[\sum_{u'} (\delta Z_{u'u}^{L*} - \delta Z_{uu'}^L) V_{u'd} + \sum_{d'} V_{ud'} (\delta Z_{d'd}^L - \delta Z_{dd'}^{L*}) \right], \quad (4)$$

where, as expected for a unitary matrix, the renormalization of V is expressed in terms of anti-hermitian matrices.

Any renormalization prescription that preserves the above WTI leads to a gauge-independent definition of the CKM matrix. To prove this theorem we start noticing that, as a consequence of $\mathcal{W}_+(\Gamma) = 0$, one has

$$\partial_\xi \mathcal{W}_+(\Gamma) = \mathcal{W}_+(\partial_\xi \Gamma) + \sum_{u,d} \left[\bar{\psi}_u^L \partial_\xi V_{ud} \frac{\vec{\delta}}{\delta \bar{\psi}_d^L} - \frac{\overleftarrow{\delta}}{\delta \psi_u^L} \partial_\xi V_{ud} \psi_d^L + \dots \right] \Gamma = 0 \quad (5)$$

where ξ is a gauge parameter and the ellipses stand for additional contributions not relevant for us. On the other hand, the gauge variation of Γ is controlled by a Slavnov-Taylor Identity (STI), $\partial_\xi \Gamma = \mathcal{S}(\partial_\chi \Gamma)$ [5, 8], where χ is the anti-commuting source of the composite operator generated by the variation of ξ , and $\mathcal{S}$ is the Slavnov-Taylor operator [9]. As $\mathcal{W}_+$ does not depend on χ and commutes with $\mathcal{S}$, the first term of Eq. (5) vanishes. The only possibility compatible with the invariance of the theory (i.e. with Eq. (2)) is then that all the parameters in the square brackets of Eq. (5) are identically zero. From this $\partial_\xi V = 0$ follows.

An interesting point about Eq. (4) is that the counterterms for physical parameters¹, the CKM matrix elements, are given in terms of conventional objects like the WFR constants. From a rigorous point of view, the distinction between physical and conventional objects can be formulated in terms of the cohomology classes of the Slavnov-Taylor operator of the theory [9]. The physical parameters on which the S -matrix depends are the coefficients of the cohomology classes. For example, in the absence of mixing, one cohomology class is provided by each Yukawa interaction term of the Lagrangian, which guarantees that the masses are physical objects. In the case of mixing, the Yukawa couplings are complex non-diagonal matrices and can be diagonalized by a redefinition of the fields, that is a finite WFR. In higher orders, however, the redefinition of fields originated by the diagonalization of the Yukawa matrix inevitably mixes with the one generated by the anomalous rescaling of the kinetic terms. A formal cohomological analysis [7] shows that indeed the off-diagonal field redefinition contains some physical parameters, namely the CKM matrix elements. On the other hand, Eqs. (2-4) allow us to disentangle the physical information related to the diagonalization of the Yukawa matrix (contained in the CKM matrix elements and in their counterterms) from the unphysical information carried by the Z factors. These constraints rely on the gauge invariance of the theory and, in a consistent framework, should be all satisfied.

As we have seen, the definition of the CKM matrix at higher orders is conveniently expressed in terms of the anti-hermitian component of the WFR. It therefore depends

¹Strictly speaking, the CKM element V_{ud} is *not* a physical quantity; physical observables are identified at tree level by the Jarlskog reparametrization invariants [10], which are constructed from physical amplitudes. This scheme can in principle be generalized at higher orders.

on the scheme chosen for the WFR. It should be clear, on the other hand, that once the counterterm δV is calculated through Eq. (4), it can be used independently of the choice of the δZ factors, because physical amplitudes are *independent* of the scheme adopted for the WFR. For example, in practical applications at the level of S -matrix, it is often convenient to avoid the rescaling of the fields (i.e. the WFR) altogether [11] and introduce only the LSZ factors for the external fields: if mixing is present, however, one still has to renormalize the mixing parameters.

A number of renormalization prescriptions for the CKM matrix are indeed possible; a first convenient option is the $\overline{\text{MS}}$ subtraction: by definition, assuming gauge invariant mass renormalization and after adjusting for the possible breaking of chiral invariance, it satisfies the WTI and the STI. Hence, as a consequence of the above theorem, it can be guaranteed to yield a gauge-independent ultraviolet pole $\delta V^{\overline{\text{MS}}}$ [12] to all orders (a proof can also be found following [13]). On the other hand, it is well-known that the decoupling of heavy particles is not manifest in the $\overline{\text{MS}}$ scheme. This means that if we work in the framework of an effective Lagrangian where the heavy fields (W boson and top quark) are integrated out, the dimension three and four operators that mix the quarks yield contributions to the amplitude which are not suppressed by the high mass scale. Moreover, as noted in [14], these terms are not even defined in the limit of vanishing light quark masses. All this makes the $\overline{\text{MS}}$ definition unnatural in the context of effective Lagrangians. As the CKM elements are mostly determined from low-energy hadronic processes, this is not very convenient. Physical amplitudes calculated with an $\overline{\text{MS}}$ counterterm for the CKM would depend on the renormalization scale (see [15] for studies of the scale evolution of the CKM matrix) and would contain $O(\alpha)$ corrections proportional to $(m_i^2 + m_j^2)/(m_i^2 - m_j^2)$, where $m_{i,j}$ are the poorly known light quark masses.

A second possibility consists in fixing four CKM elements in terms of four physical amplitudes, e.g. of the four most precise experimental processes. This procedure bypasses the definition of the WFR, but destroys the symmetry between the quark generations and is not practical in higher order calculations.

A third option is provided by the use of the *on-shell* renormalization conditions of Ref.[16] to define the WFR constants and, through Eq. (4), the CKM counterterm, as has been suggested by Denner and Sack [3] and generalized to extended models in [4]. This approach also implies decoupling in the sense explained before, i.e. dimension three and four operators are removed. The renormalization conditions, which are written in the u sector in terms of the fermionic two-point functions $\Gamma_{\bar{u}u'}(\not{p})$,

$$\begin{aligned} \bar{u}(m_u) \Gamma_{\bar{u}u'}(\not{p}) &= 0; & \Gamma_{\bar{u}u'}(\not{p}) u(m_{u'}) &= 0; \\ \bar{u}(m_u) \Gamma_{\bar{u}u}(\not{p}) \frac{1}{\not{p} - m_u} &= 1; & \frac{1}{\not{p} - m_u} \Gamma_{\bar{u}u}(\not{p}) u(m_u) &= 1, \end{aligned} \quad (6)$$

fix the masses of the u quarks and all the $\delta Z_{uu'}$ by setting them equal to the LSZ factors. Their use in Eq. (4) defines a counterterm δV_{ud} which makes physical amplitudes finite [3].

We now show explicitly that the latter procedure leads to gauge dependent amplitudes in one-loop calculations. To this end, we consider the decay of a W boson into two arbitrary quarks u and d that was studied in Ref. [3]. We conform to the notation of that paper and write the one-loop renormalized amplitude as

$$\begin{aligned} \mathcal{M}_{ud} = & V_{ud} \mathcal{M}_0 \left(1 + \delta_{vert} + \frac{\delta e}{e} - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W \right) \\ & + \mathcal{M}_0 \left(\frac{1}{2} \sum_{u'} \delta Z_{u'u}^{L*} V_{u'd} + \frac{1}{2} \sum_{d'} V_{ud'} \delta Z_{d'd}^L + \delta V_{ud} \right), \end{aligned} \quad (7)$$

where $\mathcal{M}_0 = -g/\sqrt{2} \bar{u}(m_u) \not{\epsilon}(M_W) a_- v(m_d)$, $\bar{u}$ and v are the spinors of the final-state quarks, $a_{\pm} = (1 \pm \gamma_5)/2$ are the right and left-handed projectors, and ϵ^μ is the polarization vector of the W boson.

Let us now consider the gauge dependence of the individual contributions to $\mathcal{M}_{ud}$. For our purposes, it is sufficient to consider only the ξ_W gauge parameter dependence. As the total amplitude in Eq. (7) must be gauge independent, $\partial_{\xi_W} \mathcal{M}_{ud} = 0$. Also the counterterms δe and δs_W , as defined in [11], do not depend on ξ_W at the one-loop level. On the other hand, the gauge variation of the proper vertex δ_{vert} is not trivial; it can be studied using the STI that governs the gauge dependence of the Green functions². On the mass shell, and after contracting with ϵ^μ and with the external quark spinors, one finds the following non-linear identity written in terms of 1PI Green functions

$$\partial_{\xi_W} \Gamma_{W^+ \bar{u} d} = \sum_{i=1,2} \left[\Gamma_{\chi_i \gamma^- W^+}^T \Gamma_{W^+ \bar{u} d} + \Gamma_{\chi_i \bar{u} \eta_{u'}} \Gamma_{W^+ \bar{u}' d} + \Gamma_{W^+ \bar{u} d'} \Gamma_{\chi_i \bar{\eta}_{d'} d} \right]. \quad (8)$$

Here $\gamma_\mu^\pm$ and $\eta_{u,d}$ are the sources associated to the BRST variation of $W_\mu^\pm$ and u, d fields and $\chi_{1,2}$ are the sources of the two independent composite operators generated by the variation of the two gauge fixing parameters $\xi_{W,1}$ and $\xi_{W,2}$. Following the common practice, we have set $\xi_W = \xi_{W,1} = \xi_{W,2}$.

At the one-loop level, Eq. (8) reduces to

$$\partial_{\xi_W} \Gamma_{W^+ \bar{u} d}^{(1)} = \frac{g}{\sqrt{2}} \sum_{i=1,2} \left[\Gamma_{\chi_i \gamma^- W^+}^{T(1)} V_{ud} + \sum_{u'} \Gamma_{\chi_i \bar{u} \eta_{u'}}^{(1)} V_{u'd} + \sum_{d'} V_{ud'} \Gamma_{\chi_i \bar{\eta}_{d'} d}^{(1)} \right] \not{\epsilon} a_- , \quad (9)$$

where the superscript (1) indicates that the proper functions are evaluated in the one-loop approximation and T that only the transverse component is considered. All terms are evaluated on the mass-shell of the physical fields. In a similar way, one may find

²A detailed illustration of this kind of STI [5, 8] will be given in [17].

STI for the two-point functions of the W boson and of the quarks. At one-loop level and adopting the standard tadpole renormalization, which consists in removing the whole tadpole amplitude, one finds for the W boson WFR factor

$$\partial_{\xi_W} \delta Z_W = \partial_{\xi_W} \frac{\partial}{\partial p^2} \Gamma_{W^+ W^-}^{T(1)}(p^2) \Big|_{p^2=M_W^2} = 2 \sum_{i=1,2} \Gamma_{\chi_i \gamma^- W^+}^{T(1)}(M_W^2). \quad (10)$$

The treatment of the quark two-point functions is slightly more involved: in the case of the u quarks, for instance, we decompose the unrenormalized self-energy according to

$$\Sigma_{uu'}(p) = \Sigma_{uu'}^L \not{p} a_- + \Sigma_{uu'}^R \not{p} a_+ + \Sigma_{uu'}^S (m_u a_- + m_{u'} a_+). \quad (11)$$

The one-loop STI for $\Sigma_{uu'}$ then reads

$$\partial_{\xi_W} \Sigma_{uu'}^{(1)}(p) = - \sum_{i=1,2} \left[\Gamma_{\chi_i \bar{u} \eta_{u'}}^{(1)}(\not{p}) (\not{p} - m_{u'}) + (\not{p} - m_u) \Gamma_{\chi_i \bar{\eta}_u u'}^{(1)}(\not{p}) \right]. \quad (12)$$

Splitting $\Gamma_{\chi_i \bar{u} \eta_{u'}}^{(1)}(m_u)$ into its left and right-handed components, $\Gamma_{\chi_i \bar{u} \eta_{u'}}^{L,R(1)}$, we find on the mass shell of the u quark

$$\begin{aligned} m_u \partial_{\xi_W} \left(\Sigma_{uu'}^{L,(1)} + \Sigma_{uu'}^{S,(1)} \right) &= m_{u'} \Gamma_{\chi_i \bar{u} \eta_{u'}}^{L,(1)} - m_u \Gamma_{\chi_i \bar{u} \eta_{u'}}^{R,(1)}, \\ m_u \partial_{\xi_W} \Sigma_{uu'}^{R,(1)} + m_{u'} \partial_{\xi_W} \Sigma_{uu'}^{S,(1)} &= m_{u'} \Gamma_{\chi_i \bar{u} \eta_{u'}}^{R,(1)} - m_u \Gamma_{\chi_i \bar{u} \eta_{u'}}^{L,(1)}. \end{aligned} \quad (13)$$

One can then use the above equations in the definition of the *on-shell* δZ^L (see for ex. Eq. (3.4) of Ref.[3]) and find

$$\partial_{\xi_W} \delta Z_{u'u}^{L*} = 2 \sum_{i=1,2} \Gamma_{\chi_i \bar{u} \eta_{u'}}^{R,(1)}(m_u). \quad (14)$$

An analogous result holds in the d sector.

Inserting Eqs. (10) and (14) into Eq. (9), we can write the gauge dependence of the vertex as

$$V_{ud} \partial_{\xi_W} \delta_{vert} = -\frac{1}{2} \partial_{\xi_W} \left(V_{ud} \delta Z_W + \sum_{u'} \delta Z_{u'u}^{L*} V_{u'd} + \sum_{d'} V_{ud'} \delta Z_{d'd}^L \right). \quad (15)$$

We observe that, as a consequence of the unitarity of V , in the last two terms of Eq. (15) the dependence on the CKM matrix factorizes; this is consistent with the fact that the STI in Eq. (8) would be exactly the same, were the CKM matrix diagonal, and (from a diagrammatical point of view) that the one-loop vertices involve one and only one charged quark current. The limit for massless fermions of Eq. (15) agrees with the expressions reported in [18]. Note also that all the gauge dependence of the vertex is contained in two-point function contributions. Using Eqs. (7) and (15), $\partial_{\xi_W} \mathcal{M}_{ud} = 0$

is reduced to $\partial_{\xi_w} \delta V_{ud} = 0$. In other words, we have shown that the gauge-parameter dependence introduced by the LSZ external field factors (that coincide with the *on-shell* WFR constants) is completely absorbed by the proper vertex. On the other hand, V_{ud} is a parameter of the bare Lagrangian and its renormalization condition, in the present framework, should preserve the gauge independence.

It may therefore seem surprising that the counterterm defined on the basis of the *on-shell* conditions is gauge-dependent. Indeed, an explicit one-loop calculation yields for the ξ_w dependent part of the WFR in n dimensions

$$\delta Z_{u'u}^{L,\xi_w*} = \frac{ig^2\mu^{4-n}}{2M_W^2} \sum_{d'} V_{ud'} V_{u'd'}^* \int \frac{d^n k}{(2\pi)^n} \frac{\xi_w M_W^2 - m_u^2 + m_{d'}^2}{(k^2 - \xi_w M_W^2)[(k+p)^2 - m_{d'}^2]} , \quad (16)$$

with $p^2 = m_u^2$ and $u \neq u'$; again an analogous expression holds for $\delta Z_{dd'}^L$. We see from Eq. (16) that the $1/(n-4)$ pole of the anti-hermitian part of $\delta Z^{L,\xi_w}$ is removed by GIM cancelations, ensuring that the divergence of δV_{ud} in Eq. (4) is gauge independent. On the contrary, the momentum dependence of the integrand in Eq. (16) spoils the GIM cancelations for the finite part of δV_{ud} in Eq. (4), which turns then out to be gauge dependent. We conclude that the W -decay amplitude calculated in the *on-shell* framework of Ref. [3] is gauge dependent. This can be understood by noting that the finite part of the WFR factors defined *on-shell* violates the WTI, in particular it does not satisfy Eq. (3).

A convenient and natural alternative to the prescription of [3], which maintains decoupling and enhances the symmetry among the quark generations, can be obtained by imposing the following conditions on the off-diagonal two-point functions:

$$\Gamma_{\bar{u}u'}(0) = 0 \quad ; \quad \Gamma_{\bar{u}u'}^L(0) \equiv \frac{\partial}{\partial \not{p}} \Gamma_{\bar{u}u'}(\not{p}) a_- \Big|_{\not{p}=0} = 0 , \quad (17)$$

for $u \neq u'$, and analogously for the d sector. These conditions do not fix the diagonal two-point functions. They also do not fix the off-diagonal hermitian part of the right-handed WFR, which however is finite and suppressed by light quark masses. We choose to set $\delta \mathcal{Z}_{uu'}^{R,H} = 0$ for $u \neq u'$. The normalization point, $\not{p} = 0$, is the same for all flavors and all divergences related to the mixing are correctly subtracted, as they are logarithmic and do not depend on masses and momenta. In addition, Eqs. (17) avoid problems in the treatment of the absorptive parts of the two-point functions whenever any of the quarks is heavy (typically, in the SM, this is the case with the top quark).

At the one-loop level, the antihermitian and hermitian WFR factors $\delta \mathcal{Z}_{uu'}^{L,A}$ and $\delta \mathcal{Z}_{uu'}^{L,H}$ obtained from Eq. (17) can be expressed in terms of self-energies evaluated at zero momentum transfer:

$$\delta \mathcal{Z}_{uu'}^{L,A} = \frac{m_u^2 + m_{u'}^2}{m_u^2 - m_{u'}^2} [\Sigma_{uu'}^L(0) + 2 \Sigma_{uu'}^S(0)] ; \quad \delta \mathcal{Z}_{uu'}^{L,H} = -\Sigma_{uu'}^L(0) . \quad (18)$$

It is straightforward to verify that $\delta\mathcal{Z}_{uu'}^{L,H}$ satisfies the condition of Eq. (3) because it reduces it to the case of no mixing. The CKM counterterm obtained by the use of $\delta\mathcal{Z}^{L,A}$ in Eq. (4) is, in units of $g^2/(64M_W^2\pi^2)$,

$$\begin{aligned} \delta\mathcal{V}_{ud} = & \sum_{u' \neq u} \sum_{d'} V_{u'd} V_{ud'} V_{u'd'}^* \left\{ \frac{(m_u^2 + m_{u'}^2) m_{d'}^2}{2(m_u^2 - m_{u'}^2)} \left[\frac{3}{\epsilon} - 3 \ln \frac{M_W^2}{\mu^2} \right. \right. \\ & \left. \left. - \frac{11 - 5y}{2(y-1)} - \frac{3(y-2)y}{(y-1)^2} \ln y \right] \right\} + (u \leftrightarrow d, u' \leftrightarrow d'), \end{aligned} \quad (19)$$

where $y = m_{d'}^2/M_W^2$ and $\epsilon = (4-n)/2$. $\delta\mathcal{Z}_{uu'}^{L,A}$ and $\delta\mathcal{V}_{ud}$ are gauge independent, as can be directly seen from Eq. (12). We stress once more that the use of this counterterm, based on Eq. (17) and consequently on the $\mathcal{Z}^{L,A}$ factors, is *independent* of the choice of WRF in the rest of a calculation, and corresponds to just one of the many gauge-invariant definition of the one-loop CKM elements. Needless to say, it is always the LSZ procedure (see Eqs. (6)) to dictate the treatment of the external lines. $\delta\mathcal{V}_{ud}$ can therefore be used without modifications in the calculation of [3]: in that case, the results for the W -decay amplitude are gauge independent and differ from [3] by gauge dependent $O(G_\mu m_{light}^2)$ terms.

We now consider the consistency of the renormalization conditions of Eqs. (17) beyond one-loop. First, we can show that the Eqs. (17) respect the WTI for the $W\bar{u}d$ -vertex at all orders; this WTI reads

$$\begin{aligned} (p_u + p_d)_\mu \Gamma_{\hat{W}_\mu^+ \bar{u}d}(p_u, p_d) - M_W \Gamma_{\hat{G}^+ \bar{u}d}(p_u, p_d) \\ - \frac{g}{\sqrt{2}} \left(\sum_{d'} V_{ud'} a_+ \Gamma_{\bar{d}'d}(p_d) - \sum_{u'} V_{u'd} \Gamma_{\bar{u}u'}(-p_u) a_- \right) = 0. \end{aligned} \quad (20)$$

At the one-loop level and for on-shell amplitudes, Eq. (20) holds even in the case the external Goldstone boson G^+ and the W^+ are quantized. Differentiating with respect to p_u and p_d and setting all momenta to zero, one finds that

$$\sum_{d'} V_{ud'} \Gamma_{\bar{d}'d}^L(0) - \sum_{u'} \Gamma_{\bar{u}u'}^L(0) V_{u'd} = V_{ud} G_{\bar{u}d}(0), \quad (21)$$

where $G_{\bar{u}d}(0)$ is a convergent term induced by $\Gamma_{\hat{G}^+ \bar{u}d}$. The second condition of Eqs. (17) reduces this constraint to the case of no mixing at all orders. This is the crucial requirement. Similarly, the first condition of Eqs. (17), used in Eq. (20) at zero momenta, reduces it to the well-known constraint on the renormalization of the Yukawa coupling in the absence of mixing.

We also investigate the effect of Eqs. (17) on the STI: at the one-loop level, they induce several constraints on the renormalization of χ -dependent Green functions which appear in Eq. (12), e.g. $\Gamma_{\chi i \bar{u} \eta_{u'}}^{(1)}$; using them together with Eqs. (17) in the two-loop STI,

the latter can be linearized and reduces to its one-loop form at $p = 0$. This is a non-trivial result, as the gauge dependence of the two-loop off-diagonal $\Sigma_{uu'}^{(2)}(p)$ can now be written only in terms of two-loop Green functions. At the one-loop level several simplifications occur: for example one has $\Gamma_{\chi_i \bar{u} \eta_{u'}}^{(1)}(0) a_- = 0$, because only the left-handed source $\eta_{u'}^L$ is involved. Beyond one-loop, $\eta_{u'}^R$ may also contribute, to the effect that this and analogous simplifications are not granted any longer. Consequently, the combination $\Sigma_{uu'}^L(0) + 2 \Sigma_{uu'}^S(0)$ is not guaranteed to be gauge invariant at two or more loops, although analogous but more complicated gauge-invariant combinations do exist. As already mentioned, what is certainly gauge invariant at all orders is the $\overline{\text{MS}}$ pole of δV ; we have explicitly verified this property using the STI.

A related problem concerns Σ^R , which is finite at one-loop just because of the GIM mechanism, but it may require a subtraction beyond one-loop; this would also modify the conditions of Eqs. (17). Moreover, we note that a rigorous analysis of all the WTI (not only of Eq. (20)) at higher orders cannot be performed without specifying the whole set of normalization conditions. The previous points show that a complete discussion of a non- $\overline{\text{MS}}$ renormalization of the mixing matrix beyond one-loop becomes extremely complex [17], and is only partially simplified by Eqs. (17).

The renormalization conditions of Eqs. (17) can be used in any model containing Dirac fermion mixing. For example, all has been said applies directly to the case of lepton mixing, which is suggested by recent experiments, if the neutrinos have Dirac masses. They can also be generalized to extended models, along the lines of Refs. [4].

In summary, we have reanalyzed the renormalization of the fermion mixing parameters in the Standard Model. We have reviewed several possibilities for the definition of the CKM matrix at higher orders, showing the constraints they have to satisfy in order to respect gauge invariance. In particular, we have demonstrated that the prescription based on the *on-shell* wave function renormalization constants is not consistent with the Ward-Takahashi Identities and leads to gauge dependent physical amplitudes. We have therefore proposed a simple scheme that naturally satisfies all theoretical requirements and is very convenient for practical calculations.

We thank Bernd Kniehl for suggesting the topic, for a careful reading of the manuscript and for many useful suggestions and remarks. We are also grateful to D. Maison for interesting discussions and for reading the manuscript, to A. Denner and T. Mannel for communications regarding Ref.[3], to M. Steinhauser for technical help, and to C. Becchi, G. Degrassi, and A. Sirlin for stimulating conversations.

References

- [1] W. Marciano and A. Sirlin, *Nucl. Phys.* **B93** (1975) 303.
- [2] J.F. Donoghue, *Phys. Rev.* **D19** (1979) 2772.
- [3] A. Denner and T. Sack, *Nucl. Phys.* **B347** (1990) 203.
- [4] B.A. Kniehl and A. Pilaftsis, *Nucl. Phys.* **B474** (1996) 286; S. Kiyoura, M.M. Nojiri, D.M. Pierce, Y. Yamada *Phys. Rev.* **D58** (1998) 075002.
- [5] H. Kluberg-Stern and J.B. Zuber, *Phys. Rev.* **D12** (1975) 467 and *Phys. Rev.* **D12** (1975) 482.
- [6] L.F. Abbott, *Nucl. Phys.* **B185** (1981) 189 and *Acta Phys. Pol.* 12 (1982); A. Denner, S. Dittmaier, and G. Weiglein, *Nucl. Phys.* **B440** (1995) 95.
- [7] P.A. Grassi, MPI-Th/98-57 and *Nucl. Phys.* **B462** (1996) 524.
- [8] N.K. Nielsen, *Nucl. Phys.* **B101** (1975) 173; K. Sibold and O. Piguet, *Nucl. Phys.* **B253** (1985) 517; R. Haussling and E. Kraus, *Z. Phys.* **C75** (1997) 739.
- [9] C. Becchi, A. Rouet and R. Stora, *Comm. Math. Phys.* 42 (1975) 127, *Ann. Phys.* (NY) 98 (1976) 287.
- [10] C. Jarlskog, *Phys. Rev. Lett.* **55** (1985) 1039 and *Z. Phys.* **C29** (1985) 491.
- [11] A. Sirlin, *Phys. Rev.* **D22** (1980) 971.
- [12] T.P. Cheng, E. Eichten, L.F. Li, *Phys. Rev.* **D9** (1974) 2259; M.E. Machacek, M.T. Vaughn, *Nucl. Phys.* **B222** (1983) 83.
- [13] D.J. Gross and F. Wilczek, *Phys. Rev. Lett.* **30** (1973) 1343.
- [14] P. Gambino, A. Kwiatkowski, and N. Pott, hep-ph/9810400.
- [15] K. Sasaki, *Z. Phys.* **C32** (1986) 149; K.S. Babu, *Z. Phys.* **C35** (1987) 69; C. Balzereit, T. Mannel, and B. Plümper, hep-ph/9810350.
- [16] K. Aoki, Z. Hioki, R. Kawabe, M. Konuma, T. Muta, *Prog. Theor. Phys. Suppl.* 73 (1982) 1.
- [17] P. Gambino and P.A. Grassi, in preparation.
- [18] G. Degrandi and A. Sirlin, *Nucl. Phys.* **B383** (1992) 73.